

Research Paper

Adaptation Studies of Mung bean (*Vigna radiata* L.) Varieties under Irrigated Condition in the Great Rift Valley of Ethiopia: the Case of Middle Awash Rift Valley, Werer Station.

Shimelis Alemayehu

Ethiopian Institute of Agricultural Research: Werer Agricultural Research Center: Werer/Afar, Ethiopia. P.o. Box, 2003: Addis Ababa, Ethiopia. www.eiar.gov.et. Tel: +251913087826/+251911458426.
Corresponding author's E-mail: shimitiwarc2012@gmail.com

Accepted 28 February 2022

Mung bean is one of the most important pulse crops, grown from the tropical to sub-tropical areas around the world. In Ethiopia, almost half of the land mass lies within warmer and lowland areas, with ample water resources and exploited less due to the major focus of crops production on mid to highland areas of the country receiving ample annual rainfall. In order to test the adaptability of mung bean in the lowland irrigated areas of Ethiopia, the current study was conducted at Werer agricultural research center, with the objectives to identify, select and recommend adaptable, high yielding and tolerant mung bean varieties, for irrigation production system in the lowland areas of middle awash rift valley, and at the same time assessing suitability of the area for *Vigna radiata* production. Five mung bean varieties were used in the study for two consecutive seasons (2019 and 2020). Result from the ANOVA revealed that, significant varietal differences were observed for all of the studied parameters in both seasons, except for grain filling period in the first season. The varieties Chinese, Showa-Robit and NVL-1 flowered and matured early at both growing seasons among the tested mung bean varieties. Grain yield was positively and significantly correlated with number of pods plant⁻¹, number of seeds pod⁻¹ and biomass weight, and negatively correlated with phenological characters. The maximum grain yield coupled with better number of pods plant⁻¹ and seeds pod⁻¹ was noted for the varieties N-26 and NVL-1 in the first season and Chinese and NVL-1 in the second season. Generally, mung bean varieties tested for their adaptation under this Great Rift Valley region having warmer air condition respond well and thus the area can be characterized as suitable for production of mung bean using irrigation water.

Key words: Great Rift Valley, mungbean, *Vigna radiata*, adaptation, lowland, irrigation, grain yield.

Cite this article as: Shimelis, A. (2022). Adaptation Studies of Mung bean (*Vigna radiata* L.) Varieties under Irrigated Condition in the Great Rift Valley of Ethiopia: the Case of Middle Awash Rift Valley, Werer Station. Acad. Res. J. Agri. Sci. Res. 10(1): 1-8

INTRODUCTION

Mungbean (*Vigna radiata* L. Wilczek), which is introduced recently, is an annual herb of the legume family. It has green skin and is also called green bean (MoA, 2011). It is sweet in flavor and cold in nature (EPP, 2004). The crop matures early; special features include high yield, good nutritive value, the earliness, drought resistant features and the reasonable cost of production. It is a warm season annual grain legume and the optimum temperature range for good production is 27 to 30°C and requiring 90 to 120 days of frost-free conditions from planting to maturity depending on the variety (Itefa, 2016).

Mung bean is one of the most important pulse crops, grown from the tropical to sub-tropical areas around the world (Kumari et al., 2012; Khan et al., 2012). It is an important wide spreading, herbaceous and annual legume pulse crop cultivated mostly by traditional farmers (Ali et al., 2010). Mung bean has good potential for crop rotation system, for crops under drier farmland cultivation areas (Ashraf et al., 2003) and ability of growing on dry and irrigated conditions (Rahim et al., 2010).

It is grown in several types of cultivation systems, including sole cropping, intercropping, multiple cropping and relay cropping, where it is planted after cereals using residual moisture (Rehman et al., 2009). According to Asfaw et al. (2012) in Ethiopia mung bean is mostly grown by smallholder farmers under drier marginal environmental condition and the production capacity is lower than other pulse crops. Green mung bean is less used domestically, but it is a common ingredient in Chinese and Indian cuisines. It is attributed with having high nutritional value, including protein content (24 to 26%), and helps reduce cholesterol and diabetes (Ali and Gupta, 2012; Habte, 2018).

Majority of crops production in Ethiopia is concentrated in high and mid-lands receiving rainfall. Hence, production of crop plants like mung bean using irrigation under arid and semi-arid regions of the country was kept neglected for decades. Therefore, this experiment was initiated for identifying, selecting and recommending, adaptable, high yielding and tolerant mung bean varieties, under full irrigated production system in pastoral and agro-pastoral community of the Great Rift Valley region, in the middle awash rift valley, and at the same time assessing suitability of the area for production of *Vigna radiata*.

MATERIALS AND METHODS

Description of the Study Area

The study was conducted at Werer Agricultural Research Center, found in Gebiresu Zone of Afar

Regional State, cover a long broad alluvial plain along the right bank of the Awash river. The elevation is at about 740 *masl* and located at 9°20'31" N latitude and 40°10'11" E longitude in the Middle Awash Rift Valley, close to the main high way linking Addis Ababa to Djibouti, and 260 km far from the capital of Ethiopia, Addis Ababa.

The climate is semi-arid with a bimodal rainfall of 533 mm annually. The mean minimum temperature is 15.2°C and the mean maximum temperature is 38°C. Mean relative humidity is lowest in June at 36% and highest in August at 58%. The area receives the average daily sunshine of 8.5 hours with an average solar radiation of 536 calories square centimeter⁻¹ day⁻¹ (Girma and Awulachew, 2007).

The soils are brown in color and turn to dark brown when moist. The pH of the soil is slightly alkaline and ranges from 7.5 to 8.5. The major crops grown in the area are, cotton and sugar cane with minor crops including maize, sesame, rice, wheat, pulses like chickpea and common bean, date palm, banana and vegetables in some areas around Werer Agricultural Research Center and other areas in the region.

Plant Materials and Experimental Procedures

Field evaluations of five mung bean varieties (MH-97-1, Chinese, N-26, NVL-1 and Showa-Robit) were used in the study. The particular study was conducted during the off-seasons of 2019 and 2020 at Werer Agricultural Research Center, under fully irrigated condition. The treatments were laid in RCB design with three replications having plot size of 9.6m² (4m x 2.4m), accommodating 8 rows of 4m length. The spacing between rows and plants were 30cm and 5cm, respectively. Harvesting and data collection was done from six central rows and each plot was converted to net harvestable area of 7.2m². The plots were surface irrigated by the irrigation interval of 12-15 days that varied depending on climatic and edaphic conditions.

Measurements of Crop Phonology, Plant Growth and Yield Parameters

The study consisted of crop phonology parameters (days to 50% flowering (DF), days to physiological maturity (DM) and grain filling period (GFP)); growth parameters (plant height (PH)) and yield parameters (number of pods plant⁻¹ (NPP), number of seeds pod⁻¹ (NSP), biomass yield (BMY), hundred seeds weight (HSW), harvest index (HI) and grain yield (GY)). DF and DM were recorded for each variety on the plot basis by regular observation, when 50% or more of the plants flowered, as days to 50% flowering (DF) and when 90%

of the pods in a plot dried, as days to physiological maturity (DM) of each plot. All crop phenology characters in the study were expressed in number of days; plant height in centimeters (cm); biomass yield and grain yield in kg ha^{-1} ; and weight of hundred seeds and harvest index were expressed in gram (gm) and percentage (%), respectively.

Data Collection and Statistical Analysis

At physiological maturity, five plants from central rows were randomly selected and PH in centimeters (cm) was determined. At harvest, five sample plants were randomly selected and yield components like PPP, SPP and HSW were recorded. HI (%) was calculated as ratio of GY to total BMY*100. GY was collected from six central rows of each plot (7.2m^2) and the harvested aerial plant parts were air dried at field condition for 48hrs to determine BMY. GFP, which was defined as the number of days from 50% flowering to number of days to physiological maturity was also computed and analyzed. All measured parameters (crop phenology, growth parameters, yield and yield components) were subjected to analysis of variance (ANOVA) using PROC GLM of SAS software version 9.1 (Anonymous, 2002) and the significance of mean differences were tested by least significant difference test $p \leq 0.05$ (LSD) as stated in Gomez and Gomez (1984).

Correlation Coefficient (r)

A correlation coefficient is a coefficient that illustrates a quantitative measure of some type of correlation and dependence, meaning statistical relationships between two or more random variables or observed data values. Several correlation coefficient types have been widely used. Pearson's correlation (also called Pearson's R) is a correlation coefficient that was commonly used in linear regression. Therefore, for this particular experiment, Pearson's correlation (r) was used to illustrate statistical relationships among the studied traits and calculated using the formula below.

$$r_{XY} = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2} \sqrt{\sum_{i=1}^n (Y_i - \bar{Y})^2}}$$

RESULT AND DISCUSSIONS

Crop Phenology and Growth Parameters

Days to 50% Flowering (DF)

The effect of variety on days to 50% flowering (DF), was significant ($P \leq 0.05$) in the first season and highly

significant ($P \leq 0.01$) in the second season (Table 1), in which the early days to 50% flowering were observed in variety Chinese (37.67 days) followed by Showa-Robit (38.00 days) and NVL-1 (40.33 days) on the first year (2019), whereas in the second year (2020), the variety N-26 and Chinese (36.67 days) followed by MH-97-1 (37.00 days) and Showa-Robit (39.00 days) were flowered early. Inversely, varieties MH-97-1 (43.33 days) and NVL-1 (39.67) were flowered late in both first and second year of experiments, respectively (Table 2).

Days to Maturity (DM)

The effect of variety on days to maturity (DM) was highly significant ($P \leq 0.01$) and significant ($P \leq 0.05$) both on first and second year, respectively (table 1). According to the result obtained, early matured variety was Chinese (67.33 days), followed by Showa-Robit (68.33 days) and MH-97-1 (71.00 days) on the first year, whereas in the second year, the variety Chinese (64.67 days) followed by MH-97-1 (65.00 days) and N-26 (66.67 days) matured early. Inversely, varieties NVL-1 (73.33 days) and Showa-Robit (68.00 days) were matured late in both first and second year of experiments, respectively (Table 2).

Plant Height (PH)

Result obtained from the studied traits of mung bean revealed that, plant height (PH) exerted highly significant ($P \leq 0.01$) varietal differences in both growing years. Significantly higher plant stature was observed on the variety Showa-Robit (42.54cm), followed by NVL-1 (30.63cm) and N-26 (29.93cm) during the first growing year, whereas in the second year, the variety Showa-Robit (47.33cm) followed by NVL-1 (46.54cm) and MH-97-1 (44.41cm) exhibited longest plant statures. Inversely, varieties Chinese (25.63 cm) and N-26 (40.07cm) were exhibited relatively shortest plant stature during the first and second growing periods, respectively (Table 2).

Similarly, Teame *et. al.*, (2017), reported presence of significant varietal differences among the tested varieties for crop phenological characters (DF, DM), NPP, NSP, HSW and GY. Mequannit A. and Tefera A. (2020) also reported presence of significant varietal differences among the tested varieties on GY and HSW in mungbean. In addition, Shimelis A. and Alemu D., (2019), reported presence of highly significant varietal differences for phenological characters, growth and yield components in common bean varieties studied at the same location. Biru and Dereje (2014) also reported presence of significant varietal differences for GY, DF, DM, PPP, SPP, HSW, and PH in common bean.

Table 1. Analysis of Variance for sum of squares of Mung bean traits grown under irrigation at Werer during both growing seasons (2019 and 2020).

CHARACTERS	Year 1 (2019)						Year 2 (2020)					
	Source of Variations						Source of Variations					
	Varieties (4) [‡]	Replication (2)	Error (8)	Mean	CV (%)	R ²	Varieties (4)	Replication (2)	Error (8)	Mean	CV (%)	R ²
DF	16.23*	2.47	2.38	40.07	3.85	0.79	6.10**	0.60	0.60	37.80	2.05	0.84
DM	20.77**	4.07	1.57	70.53	1.77	0.88	5.90*	2.47	1.05	66.27	1.55	0.77
GFP	12.27ns	3.47	4.47	30.47	6.94	0.61	3.27*	1.07	0.82	28.47	3.18	0.70
NPP	65.41**	10.08	4.58	20.57	10.41	0.88	31.99*	9.93	6.05	30.57	8.05	0.75
NSP	1.53**	0.59	0.14	10.44	3.61	0.87	2.84**	0.72	0.26	10.81	4.75	0.86
PH (cm)	124.36***	15.40	7.34	31.55	8.59	0.90	34.89**	8.68	3.10	43.73	4.03	0.86
BMY (kg ha ⁻¹)	9889202**	176419	1075580	8499	12.20	0.82	2305078*	2315625	452344	7138	9.42	0.79
GY (kg ha ⁻¹)	147332**	353	10317	1831	5.55	0.88	64799*	159499	14252	1930	6.19	0.84
HI (%)	23.97*	0.56	3.54	22.08	8.51	0.77	19.66**	0.34	1.16	24.32	4.43	0.90
HSW (gm)	1.12**	0.04	0.15	6.08	6.37	0.79	0.96*	0.24	0.14	5.84	6.43	0.79

Key: *, **, ns indicate significance at 0.05 and 0.01 probability levels, and non-significant, respectively. ‡=figures in parenthesis refers to degrees of freedom, CV= coefficient of variation. DF=Days to flowering, DM=Days to maturity, GFP=Grain filling period, NPP=Number of pods per plant, SPP=Number of seeds per pod, PH= Plant height, BMY=Biomass yield, YLD=Yield HI=Harvest index and HSW=Hundred seed weight

Table 2. Combined analysis of variance for sum of squares of Mung bean traits grown under irrigation at Werer during 2019 and 2020 growing seasons.

CHARACTERS	Source of Variations					
	Varieties (4) [‡]	Replication (2)	Year (1)	Variety*Year(4)	Error (18)	CV (%)
DF	35.87**	0.87	38.53**	53.47**	29.13	3.27
DM	64.20**	9.8	136.53**	42.47**	24.20	1.70
GFP	32.80*	8.27	30.00**	29.33*	43.07	5.25
NPP	346.62**	5.69	750.00**	42.99 ^{ns}	119.39	10.07
NSP	12.60**	1.17	1.00 ^{ns}	4.90**	4.71	4.82
PH (cm)	502.99**	26.02	1112.89**	133.99**	105.72	6.44
BMY (kg ha ⁻¹)	11689609*	1957148	13894681**	37087510**	15250328	11.77
YLD (kg ha ⁻¹)	233029ns	145506	72533 ^{ns}	615494**	370743	7.63
HI (%)	129.82**	0.12	37.43**	44.67**	39.26	6.37
HSW (gm)	7.93**	0.46	0.43 ^{ns}	0.39 ^{ns}	2.41	6.15

Key: *, **, ns indicate significance at 0.05 and 0.01 probability levels, and non-significant, respectively. ‡=figures in parenthesis refers to degrees of freedom, CV= coefficient of variation. DF=Days to flowering, DM=Days to maturity, GFP= Grain filling period, PPP=Number of pods per plant, SPP=Number of seeds per pod, PH= Plant height, BMY=Biomass yield, GY= Grain Yield HI=Harvest index and HSW=Hundred seed weight.

Table 3. Mean performances of Mung bean varieties for *crop phenology and growth traits*.

S/N	VARIETY	Year	DF	DM	GFP	PH (cm)
1	MH-97-1	1	43.33 ^a	71.00 ^a	27.67 ^b	28.99 ^b
		2	37.00 ^b	65.00 ^{bc}	28.00 ^b	44.41 ^a
2	Chinese	1	37.67 ^c	67.33 ^b	29.67 ^{ab}	25.63 ^b
		2	36.67 ^b	64.67 ^c	28.00 ^b	40.29 ^b
3	N-26	1	41.00 ^{ab}	72.67 ^a	31.67 ^a	29.93 ^b
		2	36.67 ^b	66.67 ^{ab}	30.00 ^a	40.07 ^b
4	NVL-1	1	40.33 ^{bc}	73.33 ^a	33.00 ^a	30.63 ^b
		2	39.67 ^a	67.00 ^a	27.33 ^b	46.54 ^a
5	Showa-Robit	1	38.00 ^c	68.33 ^b	31.67 ^a	42.54 ^a
		2	39.00 ^a	68.00 ^a	29.00 ^{ab}	47.33 ^a
GRAND MEAN		1	40.07	70.53	30.47	31.55
		2	37.8	66.27	28.47	43.73
LSD		1	2.91	2.36	3.98	5.10
		2	1.46	1.93	1.70	3.32

Key: Year1=2019, Year2=2020, DF=Days to flowering, DM=Days to maturity, GFP= Grain filling period, PH=Plant height

Table 4. Mean performances of Mung bean varieties for *Yield and Yield related components*

S/N	VARIETY	Year	NPP	NSP	BMY (kg ha ⁻¹)	GY (kg ha ⁻¹)	HI (%)	HSW (gm)
1	MH-97-1	1	16.73 ^b	10.40 ^b	8472.2 ^{bc}	1676.78 ^{cd}	19.80 ^b	5.47 ^b
		2	30.67 ^b	11.57 ^a	7250.0 ^{ab}	1887.04 ^{bc}	26.06 ^a	5.44 ^c
2	Chinese	1	17.07 ^b	9.83 ^b	6194.4 ^d	1575.31 ^d	25.67 ^a	6.82 ^a
		2	27.53 ^b	9.20 ^c	8250.0 ^a	2124.54 ^a	25.84 ^a	6.31 ^{ab}
3	N-26	1	17.73 ^b	10.19 ^b	10750.0 ^a	2121.69 ^a	19.93 ^b	6.48 ^a
		2	27.53 ^b	10.60 ^b	5937.5 ^c	1791.20 ^c	23.63 ^b	6.54 ^a
4	NVL-1	1	25.33 ^a	10.13 ^b	9722.2 ^{ab}	1979.31 ^{ab}	20.40 ^b	6.16 ^{ab}
		2	31.80 ^{ab}	11.47 ^{ab}	7562.5 ^{ab}	2039.35 ^{ab}	20.11 ^c	5.68 ^{bc}
5	Showa-Robit	1	26.00 ^a	11.67 ^a	7354.2 ^{cd}	1802.89 ^{bc}	24.62 ^a	5.44 ^b
		2	35.33 ^a	11.20 ^{ab}	6687.5 ^{bc}	1805.55 ^c	25.95 ^a	5.22 ^c
GRAND MEAN		1	20.57	10.44	8498.61	1831.2	22.08	6.075
		2	30.57	10.8	7137.5	1929.54	24.32	5.84
LSD		1	4.03	0.71	1952.7	191.24	3.54	0.73
		2	4.63	0.97	1266.3	224.77	2.03	0.71

Key: Year1=2019, Year2=2020, NPP=Number of pods per plant, SPP=Number of seeds per pod, BMY=Biomass yield, YLD=Yield HI=Harvest index and HSW=Hundred seed weight

Yield Components

Number of Pods Plant⁻¹

The productive capacity of majority of legumes is ultimately considered by the number of pods plant⁻¹. So, this makes the character one of the key factors determining the yield performance in grain legumes generally, and mung bean crop particularly.

Results obtained from the ANOVA showed that, variety had highly significant ($P \leq 0.01$) and significant ($P \leq 0.05$)

influence on number of pods plant⁻¹ during first and second growing periods, respectively (Table 1). Significantly, higher number of pods plant⁻¹ were obtained by the variety Showa-Robit (26.00, 35.33), followed by NVL-1 (25.33, 31.80) during the first and second growing seasons, respectively (Table 4). Inversely, the lowest value of the characters was exhibited by the varieties MH-97-1 (16.73) and N-26 (17.73) on the first growing season.




Figure 1. Graphic Illustration on mean of yield for the tested varieties of each growing season.




Figure 2. Graphic Illustration on mean grain yield of the tested varieties combined at both growing seasons.

Number of Seeds Pod⁻¹

Analysis of variance indicated that, there is presence of a highly significant ($P \leq 0.01$) difference among the varieties for this particular trait, both in first and second growing seasons. Better number of seeds pod⁻¹ was obtained from the varieties Showa-Robit, followed by MH-97-1 and NVL-1 in the first growing season and MH-97-1 and NVL-1 in the second growing season. Inversely, the lowest values were observed on Chinese variety during both growing seasons (Table 4). (Figure 1 and 2)

Correlation of the Characters

A Pearson correlation analysis was done in order to assess the association of various agronomic characters of mung bean in this experiment. Both positive association and non-association between characters of the component crop have been observed and discussed below.

According to the result obtained from the analysis, number of pods plant⁻¹ (0.27*), number of seeds pod⁻¹ (0.06*) and biomass yield (0.67**) affected grain yield

Table 5: Correlation coefficients (r) among 10 studied traits of mung bean varieties grown under irrigation condition at Werer (2019 and 2020).

CHARACTERS	DF	DM	GFP	NPP	NSP	PH	BMY	GY	HI
DF	1.00								
DM	0.70**	1.00							
GFP	-0.08	0.66**	1.00						
NPP	-0.41*	-0.45*	-0.19	1.00					
NSP	-0.07	-0.19	-0.20	0.48*	1.00				
PH	-0.42*	-0.57**	-0.36	0.87**	0.62**	1.00			
BMY	0.52**	0.56**	0.22	-0.29	-0.16	-0.31	1.00		
GY	-0.04	-0.03	0.01	0.27*	0.06*	0.25	0.67**	1.00	
HI	-0.74	-0.62**	-0.10**	0.35	0.02	0.30	-0.66*	0.25	1.00
HSW	-0.24	0.10	0.40*	-0.47*	-0.61	-0.55*	0.02	0.01	0.01

positively and significantly. This shows that, these factors were responsible for the production of grain yield in mung bean. This study is in line with the study by Teame *et. al.*, (2017), reported positive and significant correlation of grain yield with NPP, NSP and HSW in mung bean. In contrast, both days to flowering (DF) and days to maturity (DM) affected grain yield negatively and the observed association was very weak and non-significant (Table 5).

CONCLUSION AND RECOMMENDATIONS

Mung bean (*Vigna radiata* L. Wilczek) is an essential short duration, self-pollinated diploid legume crop with high nutritive values and nitrogen fixing ability. It is an eco-friendly food grain leguminous crop of dry land agriculture with rich source of proteins, vitamins, and minerals. As almost half of the land mass in Ethiopia lies within warmer and lowland areas, with ample water resources and exploited less due to the major focus of crops production on other areas receiving sufficient annual rainfall, there must be extension of crops adapting and responding better in the lowland areas of the country.

From this particular study, almost all of the studied characters exhibited significant and highly significant varietal differences at both growing periods except GFP during the first (2018) growing period indicating the existence of variability among the tested varieties for the studied parameters in this experimentation.

The middle awash area, in the Great Rift Valley region of Ethiopia found to be suitable agro-ecological zone for the production of mung bean. However, additional investigations has to be conducted on other crop management packages necessary for the escalation of grain yield in mung bean grown under irrigation condition of the study area, and other areas having similar agro-ecology. It can also be noted that, extension works on

these newly adapted crop varieties should be conducted to address the end producers with the varieties.

REFERENCES

- Ali M, Gupta S (2012). Carrying capacity of Indian agriculture: pulse crops. *Current Science* 102:874-881.
- Ali M.Z., Khan M.A.A., Rahaman A.K.M.M., Ahmed M. and Ahsan A.F.M.S. 2010. Study on seed quality and performance of some mung bean varieties in Bangladesh. *Int. J. Expt. Agric.* 1 (2), 10-15.
- Anonymous. 2002. SAS Guide for personal computers, version 9.0 editions. SAS Institute Inc., Cary, NC, USA.
- Asfaw A, Gurum F, Alemayehu F, Rezene Y (2012). Analysis of multi environment grain yield trials in mungbean (*Vigna radiata* L.) Wilczek based on GGE biplot in Southern Ethiopia. *Journal of Agricultural Science Technology* 14:389-398.
- Ashraf M., Mueen U.D.M. and Warraich N.H. 2003. Production efficiency of mungbean (*Vigna radiata* L.) as affected by seed inoculation and NPK application. *Int. J. Agri. Biol.* 5 (2).
- Biru A. and Dereje A. (2014). Adaptation Study of Improved Haricot Beans (*Phaseolus Vulgaris* L.) Varieties at Western Oromia, HaroSabu, Ethiopia.
- EPP (Ethiopian Pulses Profile), "Ethiopian export promotion agency, product development & market research directorate, May 2004 Addis Ababa, Ethiopia," 2004.
- Girma, M.M. and Awulachew, S.B. (2007). Irrigation practices in Ethiopia: Characteristics of selected irrigation schemes. Colombo, Sri Lanka: International Water Management Institute, 80p. (IWMI Working Paper 124).
- Girma, M.M. and Awulachew, S.B. (2007). Irrigation practices in Ethiopia: Characteristics of selected

- irrigation schemes. Colombo, Sri Lanka: International Water Management Institute, 80p. (IWMI Working Paper 124).
- Gomez, K. A. and A. A. Gomez. (1984). Statistical Procedures for Agricultural Research. 2nd Ed., John Willey and Sons. Singapore
- Habte U (2018). Evaluation of adaptability of mungbean varieties in moisture stress of Eastern Harerghe Zone. *Agricultural Research and Technology* 13(2): 555880.
- Itefa D (2016). General characteristics and genetic improvement status of mungbean (*Vigna radiata* L.) in Ethiopia: Review article. *International Journal of Agriculture Innovations and Research* 5(2):2319-1473.
- Khan M.A., Naveed K., Ali K., Ahmad B. and Jan S. 2012. Impact of mung bean-maize intercropping on growth and yield of mung bean. *Weed science society of Pakistan department of weed science. J. Weed Sci. Res.* 18 (2): 191-200.
- Kumari R., Shekhawat K.S., Gupta R. and Khokhar M.K. 2012. Integrated management against root- rot of mung bean (*Vigna radiata* (L.) Wilczek) incited by *macrophomina phaseolina*. *J Plant Pathol Microb* 3:5.
- Mequannit Aklilu and Tefera Abebe, 2020. Adaptation study of mung bean (*Vigna radiata*) varieties in tepi, south western Ethiopia, Pelagria Research Library, *Asian Journal of Plant Science and Research*, 10(5):58-61.
- MoA (2011). Ministry of Agriculture: Animal and Plant Health Regulatory Directorate. Crop Variety Register Issue No. 14. Addis Ababa, Ethiopia.
- Rahim M.A., Mia A.A., Mahmud F., Zeba N. and Afrin K.S. 2010. Genetic variability, character association and genetic divergence in mungbean (*Vigna radiata* L. Wilczek). *POJ* 3 (1):1-6.
- Rehman A., Khalil S.K., Nigar S., Rehman S., Haq I., Akhtar S., Khan A.Z. and Shah S.R.. 2009. Phenology, plant height and yield of mungbean varieties in response to planting date. *Sarhad J. Agric.* 25 (2): 147-151.
- Shimelis, A., Alemu, D. (2019). Adaptation Study of Improved Common Bean (*Phaseolus vulgaris* L.) Varieties under Irrigated Condition in the middle Awash Rift Valley, Werer, Ethiopia. *Acad. Res. J. Agri. Sci. Res.* 7(1): 1-7
- Team G., and Bisrat G., (2017). Adaptation Study of Mung Bean (*Vigna Radiata*) Varieties in Raya Valley, Northern Ethiopia. *Current Research in Agricultural Sciences, Vol. 4, No. 4, pp. 91-95.*